



# Corporate Presentation

January 2021



SEEKING GENE THERAPY CURES

NASDAQ: RCKT

# Important Information

---

## Cautionary Statement Regarding Forward-Looking Statements

Various statements in this release concerning Rocket's future expectations, plans and prospects, including without limitation, Rocket's expectations regarding its guidance for 2020 in light of COVID-19, the safety, effectiveness and timing of product candidates that Rocket may develop, to treat Fanconi Anemia (FA), Leukocyte Adhesion Deficiency-I (LAD-I), Pyruvate Kinase Deficiency (PKD), Infantile Malignant Osteopetrosis (IMO) and Danon Disease, and the safety, effectiveness and timing of related pre-clinical studies and clinical trials, may constitute forward-looking statements for the purposes of the safe harbor provisions under the Private Securities Litigation Reform Act of 1995 and other federal securities laws and are subject to substantial risks, uncertainties and assumptions. You should not place reliance on these forward-looking statements, which often include words such as "believe," "expect," "anticipate," "intend," "plan," "will give," "estimate," "seek," "will," "may," "suggest" or similar terms, variations of such terms or the negative of those terms. Although Rocket believes that the expectations reflected in the forward-looking statements are reasonable, Rocket cannot guarantee such outcomes. Actual results may differ materially from those indicated by these forward-looking statements as a result of various important factors, including, without limitation, Rocket's ability to monitor the impact of COVID-19 on its business operations and take steps to ensure the safety of patients, families and employees, the interest from patients and families for participation in each of Rocket's ongoing trials, our expectations regarding when clinical trial sites will resume normal business operations, our expectations regarding the delays and impact of COVID-19 on clinical sites, patient enrollment, trial timelines and data readouts, our expectations regarding our drug supply for our ongoing and anticipated trials, actions of regulatory agencies, which may affect the initiation, timing and progress of pre-clinical studies and clinical trials of its product candidates, Rocket's dependence on third parties for development, manufacture, marketing, sales and distribution of product candidates, the outcome of litigation, and unexpected expenditures, as well as those risks more fully discussed in the section entitled "Risk Factors" in Rocket's Quarterly Report on Form 10-Q for the quarter ended September 30, 2020, filed November 6, 2020 with the SEC. Accordingly, you should not place undue reliance on these forward-looking statements. All such statements speak only as of the date made, and Rocket undertakes no obligation to update or revise publicly any forward-looking statements, whether as a result of new information, future events or otherwise.

# Mission, Vision and Values

## TRUST



Trust is given and trust is earned – it's a balance. The word trust comes from the Proto-Indo-European word deru which means "to be firm, solid, steadfast." Trust is the ground and foundation for everything we do.

## GENEROSITY



Being generous means following up, sharing our best ideas, forgiving ourselves and others, asking who needs us, treating our word as gold, taking time to truly see others, and so many other things. The word generous has the same root as the word "gene" – which meant "to beget." Genes thrive on the generosity of others. What more is there to say?

## CURIOSITY




The wonder of a child staring up at the night sky. Humility, egolessness. No single one of us can do this job alone and it is ok to ask for help. Curiosity is derived from the Latin word "cura" which gave birth to the word "care" as well as "cure." Generosity is to curiosity what gene is to cure.

## ELEVATE




Derived from Latin levis which means "light" as opposed to heavy. How can we bring trust, generosity and curiosity to elevate ourselves, each other, the pipeline and ultimately the life experience of patients and their families?

# Multi-Platform Gene Therapy Targeting Rare Diseases




# Gene Therapy: A Multi-Platform Approach

## In Vivo (In Body) Adeno-associated Virus Gene Therapy



## Ex Vivo (Outside Body) Lentiviral Gene Therapy



# About Rocket Pharma

**Multi-Platform Gene Therapy  
Company Targeting Rare Diseases:  
1st-in-class with direct on-target  
mechanism of action and clearly-  
defined clinical endpoints**

**Ex-vivo Lentiviral vectors (LVV)**

- Fanconi Anemia (FA)
- Leukocyte Adhesion Deficiency-I (LAD-I)
- Pyruvate Kinase Deficiency (PKD)
- Infantile Malignant Osteopetrosis (IMO)

**In-vivo adeno-associated virus (AAV)**

- Danon Disease

**Multiple Near- & Medium-  
term Company Value Drivers**

**Near-term Milestones (2020)**

- All five programs in the clinic (initiation of IMO)
- New preliminary data in Danon & PKD; Additional mature data in FA & LAD-I
- Two programs in registration-enabling Phase 2 (FA, LAD-I)

**Medium-term Milestones (2021-2022)**

- First global submission (BLA)
- Platform establishment and pipeline expansion
- Current programs eligible for Pediatric Priority Review Vouchers

**Strong Precedents and  
World-Class Expertise**

**Strong Precedents and Sound Strategy**

- Compelling clinical proof-of-concept for LVV- & AAV-based therapies across a spectrum of genetic disorders
- Clearly-defined product metrics across indications
- Experienced company leadership
- Leading research and manufacturing partners

# Rocket's Leadership Team



**Gaurav Shah, M.D.**  
President & Chief Executive Officer  
Spearheaded Kymriah (CART-19)  
development at Novartis towards approval



**Kinnari Patel, Pharm.D., MBA**  
COO & Head of Development, EVP  
Led Opdivo and six rare disease indication  
approvals



**Claudine Prowse, Ph.D.**  
SVP, Strategy & Corporate Dev  
~20 years capital markets, strategy,  
corporate development



**Jonathan Schwartz, M.D.**  
CMO & Clinical Development, SVP  
Led multiple biologics approvals



**John Militello, CPA**  
VP, Principal Accounting Officer  
~20 years public company finance and accounting  
experience, 6 years biotech experience



**Raj Prabhakar, MBA**  
Chief Business Officer, SVP  
~17 years cell, gene and biotech  
business development



**Gayatri R. Rao, M.D., J.D.**  
VP, Global Program Head, LVV  
7-Year Former Director of FDA's Office of Orphan  
Products Development



**Carlos Garcia-Parada, MBA**  
Chief Financial Officer  
14 years of Oncology & Rare Disease experience  
Leading role in launching Kymriah, the first CAR-T product on the market.



**Ramji Krishnan, Ph.D.**  
VP, Manufacturing & Manufacturing Sciences  
17+ years of CMC product development and life cycle  
management expertise




**José Trevejo**  
Chief Development Officer of AAV, SVP  
~20 years of clinical development expertise



**Brian C. Beard, Ph.D.**  
AVP, CMC  
15+ years cell and gene therapies expertise



# Rocket's Expanding Pipeline: Potential for Significant Value Creation Near and Long Term



# Fanconi Anemia (FA)

## Monogenic DNA-repair disorder

RP-L102

Fanconi Anemia

RP-A501

Danon Disease

RP-L201

Leukocyte Adhesion Deficiency-I


RP-L301

Pyruvate Kinase Deficiency

RP-L401

Infantile Malignant Osteopetrosis

### OVERVIEW:



**Current available treatments:** Allogeneic hematopoietic stem cell transplant associated with 100-day mortality, GVHD, and additional increased cancer risk



**Addressable Market<sup>2</sup>:** Estimated US + Europe target population of approximately 4,000 patients, 500 patients/year



**RP-L102:** LVV gene therapy that elicits phenotypic correction of blood cells and stabilization of previously declining blood counts



**Regulatory Designations:** Fast Track, Regenerative Medicine Advanced Therapy (RMAT) and Rare Pediatric Disease designations in the US; Advanced Therapy Medicinal Product (ATMP) classification and PRiority MEDicines (PRIME) in the EU; Orphan Drug designation in the US/EU


<sup>1</sup> Alter Br J Haematol 2010.

<sup>2</sup> 4,000 based on a detailed population analysis of FA genomic variants. 500 per year extrapolated by actual transplants per year plus patients from prevalence

# Potential to Correct Bone Marrow Defect without Conditioning to Prevent Hematologic Failure


## Rationale for GTx in FA:

- Somatic mosaicism demonstrates that a **modest number** of gene-corrected hematopoietic stem cells can repopulate a patient's blood and bone marrow with corrected (non-FA) cells.<sup>1,2</sup>



## Gene Therapy Value Proposition:

- Potential to **correct** blood & bone marrow defect **without conditioning**
- GTx implemented as preventative measure to **avert bone marrow failure**; BMT is indicated for patients in whom marrow failure has occurred.



<sup>1</sup> Soulier, J., et al. (2005) Detection of somatic mosaicism and classification of Fanconi anemia patients by analysis of the FA/BRCA pathway. *Blood* 105: 1329-1336; <sup>2</sup>Data on file: Showing a single patient with a spontaneous correction of blood counts, no therapy administered.

# FA Path to Product Registration

## CIEMAT-Sponsored FANCOLEN 1 Study Process A

- Interim data (>12-month follow-up) showed evidence of durable engraftment, continued improvement in phenotypic markers and stabilization of previously-declining blood counts
- No conditioning required

## OPTIMIZATION

## Rocket-Sponsored Process B





*(Optimized CD34 cell enrichment,  
transduction enhancers,  
commercial-grade vector and  
modified cell processing)*

- Clinical trial of ~12 patients with sites at Stanford (US), Niño Jesús Hospital (Spain), and other leading centers in the US/Europe
- No conditioning required

BLA/  
MAA

# Bone Marrow Engraftment: Increasing Blood Cell VCNs Provide Evidence of Survival Advantage of Gene-Corrected FA Cells

First Demonstration of Engraftment Without Conditioning  
 (“Process A” —non-optimized—RP-L102)





\* This point requires additional validation as the long-term follow-up study is activated

HIUNJ Data Presented at ASGCT By CIEMAT May 2020  
 cCFU = Corrected Colony Forming Units; pVCN: Product VCN \*Minimally Acceptable Dose







# Functional Correction of Bone Marrow

## Progressive Phenotypic Correction of BM Cells (MMC-Resistance)




- MMC assay *identifies cells resistant* to Mitomycin-C (MMC), a DNA damaging agent toxic to (uncorrected) FA blood and bone marrow cells

# Gene Therapy Confers a Phenotype Similar to Mosaicism






# Increases of Corrected Leukocytes Support Restoration of Normal Bone Marrow Function Consistent with Mosaic Phenotype

## Kinetics of Corrected and Uncorrected PB Leukocytes Prior to and After Gene Therapy






# Gene Therapy Improves Previously Declining Blood Counts in Optimally Treated Patients

**02002 (Cryo)**  
 $2.5 \times 10^5$  cCD34<sup>+</sup>/Kg  
 $1.7 \times 10^4$  cCFU/Kg



**02006 (Fresh)**  
 $4.0 \times 10^5$  cCD34<sup>+</sup>/Kg  
 $1.6 \times 10^5$  cCFU/Kg



# RP-L102 “Process B”: Pivotal Clinical Trials and Outcome Measures

| RP-L102 Studies    | Non-randomized, open label studies: US Phase 1, US Phase 2, and EU Phase 2 (FANCOLEN-II)                                                                                                                                                                                                                                                                                                                                                                                            |                                                                                                                                                                     |
|--------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| CMC/Drug Product   | “ <b>Process B</b> ” includes cell enrichment, transduction enhancers, commercial-grade vector and modified cell processing                                                                                                                                                                                                                                                                                                                                                         |                                                                                                                                                                     |
| Inclusion Criteria | <p>Focus on patients with no/limited marrow failure, optimize preventative potential in absence of conditioning</p> <p>Minimum age: 1; Maximum age: US Ph 1 (12-yrs); US Ph 2 (none); EU Ph 2 (17-yrs)</p> <p>BM CD34+ concentration ≥ 30/μL (from aspirate); if BM CD34+ of 10-29/μL, then at least 2 of the following: Hb ≥ 11g/dL, ANC ≥ 900/μL, or Platelets ≥ 60,000/μL</p> <p>US Ph 1 only: At least 1 hematologic parameter (Hb, ANC or Plt) below lower limit of normal</p> |                                                                                                                                                                     |
| Exclusion Criteria | <p>Available &amp; eligible HLA-identical sibling donor</p> <p>MDS or leukemia (including associated cytogenetic abnormalities)</p> <p>Mosaicism with stable/improved blood counts</p>                                                                                                                                                                                                                                                                                              |                                                                                                                                                                     |
| Endpoints          | <b>Efficacy</b><br><br><b>Engraftment:</b> Peripheral blood (PB) and BM vector copy number (VCN)<br><b>Phenotypic correction:</b> Increased resistance of BM and PB cells to MMC and DEB<br><b>Clinical response:</b> Prevention of BMF                                                                                                                                                                                                                                             | <div> <b>Efficacy in 5 of 12 Patients (observed over 1-3 years post rx) required to reject null hypothesis</b> </div> <div> <b>Safety</b> of RP-L102         </div> |

# RP-L102 Treated Study Patients

| Phase   | Patient # | Site  | Age at Enrollment | Gender | Follow-up |
|---------|-----------|-------|-------------------|--------|-----------|
| PHASE 1 | 1 (1001)  | US    | 5                 | F      | 18M       |
|         | 2 (1002)  | US    | 6                 | F      | 18M       |
|         | 3 (2004)  | Spain | 3                 | M      | 12M       |
| PHASE 2 | 4         | Spain | 2                 | F      | 2M        |
|         | 5         | Spain | 3                 | M      | 2M        |
|         | 6         | US    | 3                 | M      | 2M        |
|         | 7         | US    | 5                 | F      | 2M        |
|         | 8         | UK    | 6                 | F      | 1M        |
|         | 9         | US    | 2                 | M      | -         |

- 9 patients treated across 3 clinical sites, 2 under Phase 1 and 7 under global Phase 2
- All patients  $\leq 6$ -years at enrollment
- 3 patients have  $\geq 12$ -months of follow-up; remaining treated more recently with limited follow-up
- Note: Follow-up and patient enrollment has been complicated by COVID-19 pandemic

# RP-L102 Investigational Product Metrics

| Phase   | Patient # | Follow-up | CD34+ Cells/kg <sup>^</sup> | CFCs/kg <sup>^</sup>  | Mean VCN: Liquid Culture | Mean VCN: CFCs | Transduction Efficiency (%) | CFC Survival MMC 10nM (%) |
|---------|-----------|-----------|-----------------------------|-----------------------|--------------------------|----------------|-----------------------------|---------------------------|
| PHASE 1 | 1 (1001)  | 18M       | 2.0 x 10 <sup>5</sup>       | 5.2 x 10 <sup>4</sup> | 2.08                     | 0.62           | 67                          | 33                        |
|         | 2 (1002)  | 18M       | 3.7 x 10 <sup>5</sup>       | 5.0 x 10 <sup>4</sup> | 2.21                     | 0.92*          | 72                          | 47                        |
| PHASE 2 | 3 (2004)  | 12M       | 4.8 x 10 <sup>5</sup>       | 1.1 x 10 <sup>5</sup> | 1.70                     | 0.73           | 100                         | 63                        |
|         | 4         | 2M        | 3.2 x 10 <sup>6</sup>       | 2.8 x 10 <sup>5</sup> | 1.65                     | 1.56           | 97                          | 62                        |
|         | 5         | 2M        | 1.9 x 10 <sup>6</sup>       | 1.5 x 10 <sup>5</sup> | 2.16                     | 0.76           | 61                          | 45                        |
|         | 6         | 2M        | 4.1 x 10 <sup>6</sup>       | Pending               | 0.62                     | Pending        | Pending                     | Pending                   |
|         | 7         | 2M        | 2.8 x 10 <sup>6</sup>       | Pending               | 1.46                     | Pending        | Pending                     | Pending                   |

Overall DP metrics were consistent with the more optimally treated patients from FANCOLEN-I study

## Median values:

VCN (liq) 1.7  
VCN (CFC) 0.76  
TD efficiency 72%  
CFC MMC-res 47%

Overall transduction and MMC-resistance levels in DP were consistent with high degree of corrected HSPCs

\* Mean CFC VCN was assessed from a cryopreserved drug product sample.

<sup>^</sup> Per NC200 automated count (results in ~50% lower count vs. manual used in FANCOLEN-I).

CFCs: colony forming cells

VCN: vector copy number





MMC: mitomycin-C

Data as of October 2020

Optimal means of the nine patients treated in Fancolen-I the two that had the best benefit risk







# RP-L102 Treated Study Patients (>12M Follow-up)







1001

- Prior to gene therapy, decreases in multiple blood lineages over prior 12-24 months



2004

- Post gene therapy, stability across multiple blood lineages observed over 12-18 months




1002

- One patient with *Influenza B* infection 9-months post-rx; Required transfusions and subsequent BMT at 18-months post-rx

# RP-L102 Treated Study Patients (>12M Follow-up)


- N = 3 with ≥ 12-Months of Follow-up
- 2 of 3 showed increasing evidence of engraftment
  - 1 patient's course (1002) complicated by *Influenza B* infection; received BMT
  - *Of note: Additional 3 of 4 patients with 2-months follow-up have early evidence of engraftment (0.01-0.02)*



LOQ = Limit of quantitation    BQL= Below quantitation limits    NDA = No detectable amplification    \* Early time points had gene marking that was below quantification limits (BQL)

# RP-L102 Treated Study Patients (>12M Follow-up)

**“Process B” BM VCN was in Line or Better than Optimally Treated “Process A” Patients at 12-Months**




Optimal means of the nine patients treated in Fancolen-I the two that had the best benefit risk


# RP-L102 Treated Study Patients (>12M Follow-up)

## Increasing Proportion of Gene-Corrected Cells Observed in Peripheral Blood

Patient 1001



Patient 2004



● Corrected WBC    ■ Uncorrected WBC

# Summary of Pivotal RP-L102 Treated Study Patients

**PB VCN available for N = 7**  
**5 of 7 showed preliminary evidence of engraftment**

**N = 3 with  $\geq 12M$  VCN**

- 2 of 3 showed increasing evidence of engraftment
- 1 patient's course (1002) complicated by *Influenza B* infection; required BMT

**N = 4 with early VCN (2M)**

3 showed early evidence of engraftment  
(0.01-0.02)

- All patients clinically stable post-treatment; the patient who required BMT underwent transplant at 18-months and engrafted without complications
- RP-L102 related SAEs: 1 transient infusion-related reaction (Grade 2)
- Patient enrollment and follow-up has been challenged by COVID-19 pandemic

# RP-L102 Conclusions: Optimized “Process B” Appears to be a Consistent and Reproducible Improvement over “Process A”

---

- **9 out of 12 planned patients treated** with “Process B”
  - 7 patients with follow-up data: 3 with  $\geq 12$ M follow-up
- Safety results appear **highly favorable**
  - Patients treated without conditioning
  - No signs of dysplasia or other concerning features
- Evidence of **preliminary engraftment** observed in 5 out of 7 patients to-date
  - 1 patient’s course complicated by Influenza B resulting in progressing BMF; successfully received BMT at 18-months
  - 1 patient awaiting further follow-up
- Evidence of increasing engraftment, MMC-resistance and **stable blood counts** in 2 out 3 patients with  $\geq 12$ M follow-up

**\* Efficacy activity in 5 of 12 patients (observed over 1-3 years post rx) required to reject null hypothesis**

# Danon Disease

## *Monogenic Heart Failure Syndrome*

RP-L102  
Fanconi Anemia

RP-A501  
Danon Disease

RP-L201  
Leukocyte Adhesion Deficiency-I


RP-L301  
Pyruvate Kinase Deficiency

RP-L401  
Infantile Malignant Osteopetrosis

### OVERVIEW:

-  **Background:** Devastating *multisystemic disorder* caused by highly penetrant and X-linked dominant *LAMP2 mutations*, rapidly progressive cardiomyopathy is predominant cause of morbidity and early mortality in adolescents & young adults
-  **Currently available treatments:** *Non-curative* heart transplants associated with considerable morbidity and mortality
-  **Addressable Market:** Estimated US + Europe prevalence of *15,000-30,000*
-  **RP-A501:** AAV9 gene therapy product that elicits *improvements* in *survival*, cardiac function, and liver enzymes in preclinical studies
-  **Regulatory Designations:** Orphan Drug, Rare Pediatric & Fast Track designations in the US

# An Impairment in Autophagy Caused by *LAMP2B* Mutations




# Epidemiology and Market Opportunity

## Hypertrophic Cardiomyopathy (HCM)

- US HCM Prevalence: 600K-1MM+\*
- 1-4% of HCM patients consistently identified with LAMP2 mutations in multiple studies with >1000 subjects evaluated\*\*
- Danon Disease Patients with HCM: \*\*\*
  - 85% of males
  - 30% of females

## Dilated Cardiomyopathy (DCM)

- Danon Disease Patients with DCM \*\*\*
  - 15% of males
  - 50% of females



\* J Am Coll Cardiol. 2015 Mar 31;65(12):1249-1254.

\*\* Heart. 2004 Aug;90(8):842-6. N Engl J Med. 2005 Jan 27;352(4):362-72. Genet Med. 2015 Nov;17(11):880-8. Gene. 2016 Feb 15;577(2):227-35. J Cardiovasc Transl Res. 2017 Feb;10(1):35-46

\*\*\* Neurology. 2002 Jun 25;58(12):1773-8. Genet Med. 2011 Jun;13(6):563-8. Rev Esp Cardiol (Engl Ed). 2018 Aug 11.


# Danon Disease Causes 1-4% of Hypertrophic Cardiomyopathy: *Consistent Presence in Multiple Series Published 2004-Present*

| Author & Year | Age    | n HCM | n Danon | % Danon | Note                                                                                                                                                                                                                                                  |
|---------------|--------|-------|---------|---------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Charron 2004  | N.A.   | 197   | 2       | 1.0%    | Studied LAMP2 mutations in 197 HCM patients at a general hospital in Paris                                                                                                                                                                            |
| Arad 2005     | 12-75  | 75    | 2       | 2.7%    | Studied glycogen storage diseases in 75 consecutive pts diagnosed with HCM (multicenter US/EU). No cases of Pompe or Fabry were detected.                                                                                                             |
| Yang 2005     | 1m-15y | 50    | 2       | 4.0%    | Studied LAMP2 mutations in 50 pts with ped./juvenile onset HCM (single US center). Additional DD identified in relatives of the n=2 probands were not included in this analysis.                                                                      |
| Cheng 2012    | N.A.   | 50    | 3       | 2.3%    | Studied LAMP2 mutations in 50 consecutive pts diagnosed with concentric LVH at a general hospital in Peking. (Concentric LVH is seen in appx. 38% of HCM). DD incidence higher (3/36) when n=14 w/ cardiac amyloidosis were removed from n=50 cohort. |

Charon et al. Heart 2004; 90:842-6.  
 Arad et al. N Engl J Med 2005; 352:362-72.  
 Yang et al. Circulation 2005; 112:1612-17.  
 Cheng et al. Eur Heart J 2012; 33:649-56.

# RP-A501 Restores Cardiac Function in KO Mice


## Dose-Dependent Improvements in Systolic and Diastolic Function in LAMP2 KO Mice



Lower dP/dt max indicates impaired contractility; Higher (less negative) dP/dt min indicates impaired heart relaxation

\*PBS = Phosphate Buffered Saline (Negative Control)


# RP-A501 Shows Survival Benefit at Higher Doses in Preclinical Studies




Note: All mice were sacrificed at Month 10

# Protein: RP-A501 Elicits Durable Expression of LAMP2B Protein and Autophagy in Heart<sup>1</sup>


## LAMP2 PROTEIN EXPRESSION



## LC3-II PROTEIN EXPRESSION




## WESTERN BLOT




<sup>1</sup>Data are Mean  $\pm$  SEM. N=5-8 per group. Untx = Untreated, PBS = Phosphate buffered saline  
Note: Mouse LAMP2 and Human LAMP2 data are from separate Western blots.

# Structural: RP-A501 Reduces Autophagic Vacuoles in All KO Mouse Models



# Dose-dependent LAMP2 Expression in Cardiac Tissue



# AAV9 Vector Shows Consistent Cardiac Tropism in Several Studies Across Different Species

| DISORDER & VECTOR        | DOSE                      | SPECIES     | RESULTS                                                                                | SPONSOR               | REFERENCE                |
|--------------------------|---------------------------|-------------|----------------------------------------------------------------------------------------|-----------------------|--------------------------|
| LGMD2A<br>AAV9.hCAPN3    | 3E+13 vg/kg               | NHP         | 8-80-fold higher transduction in cardiac vs. skeletal muscle                           | Genethon              | Lostal (ASGCT 2018)      |
| Non-specific<br>AAV9.Luc | 3E+12 vg/kg               | NHP         | ~ 10-fold higher transduction in cardiac vs. diaphragm; and comparable to other muscle | UNC                   | Tarantal 2016            |
| Pompe<br>AAV9.hGAA       | 1E+11 vg/mouse            | Mouse       | ~ 10-fold higher transduction in cardiac vs. diaphragm                                 | U. Florida            | Falk 2015                |
| DMD<br>AAV9.mDys         | 1.9 - 6.2E+14 vg/kg       | Dog         | 2-3 fold higher transduction in cardiac vs. skeletal muscle                            | U. Missouri           | Yue 2015                 |
| SMA<br>AAV9.SMN          | 3E+14 vg/kg & 1E+13 vg/kg | Mouse & NHP | ~ 100-fold higher transduction in cardiac vs. skeletal muscle (mouse)                  | Nationwide Children's | Meyer 2014               |
| MPSIIIB<br>AAV9.hNAGLU   | 1 - 2E+13 vg/kg           | NHP         | ≥ 10-fold higher transduction in cardiac vs. skeletal muscle in majority of animals    | Nationwide Children's | Murrey 2014              |
| Non-specific<br>AAV9.Luc | 5E+10 vg/mouse            | Mouse       | 5-10-fold higher transduction in cardiac vs. skeletal muscle                           | UNC                   | Pulicherla 2011          |
| Pompe<br>AAV9.hGAA       | 4E+05 - 4E+08 vg/mouse    | Mouse       | ~ 8-12-fold higher transduction in cardiac vs. skeletal muscle or diaphragm            | U. Florida            | Pacak 2006               |
| SMA<br>AAV9.SMN          | 2E14 vg/kg                | Human       | Heart VCN ~3-4, Muscle & CNS ~1                                                        | AveXis                | Kaspar 2019 (ASGCT 2019) |

# Summary of Preclinical Data

---

- Shows Phenotypic Improvements at Low-Dose 5e13 vg/kg:
  - **Survival** benefit at higher doses
  - Dose-dependent **restoration** of cardiac function
  - Improvement in transaminases
- RP-A501 **Reduces** Autophagic Vacuoles in All Examined Organs: Heart, Liver, Skeletal Muscle
- RP-A501 Elicits **dose-dependent increase** in LAMP2 mRNA and protein
- RP-A501 Preclinical Safety, Tox and Biodistribution Summary:
  - No therapy-related deaths
  - No significant hematologic changes
  - No significant biochemical changes
  - No significant clinical chemistry changes
  - Mild and transient ALT elevation that self-resolved after one week in a single NHP
  - In both mouse and NHPs, VCN detection in Danon disease organs indicated high *LAMP2B* presence in heart tissue (for NHP, ~10x higher on average than in skeletal muscle and CNS)

# RP-A501 Clinical Trial and Outcome Measures

## Non- Randomized Dose-Escalation Phase 1 Study

### Study Design

- Phase 1 open label study in male Danon patients
- Two age cohorts
  - Adolescent/Adult (>15 y)
  - Pediatric (8-14 y)
- Treatment doses
  - Low  $6.7 \times 10^{13}$  GC/kg
  - Higher  $1.1 \times 10^{14}$  GC/kg<sup>1</sup>

### Primary Outcomes

- Assessment of:
  - Safety at all doses
  - Target tissue transduction & LAMP2B expression
  - Effect on cardiomyocyte histology
  - Clinical stabilization or improvement via cardiac imaging, serology and exercise testing

1. IND enables doses up to  $2.0 \times 10^{14}$  GC/kg  
<https://clinicaltrials.gov/ct2/show/NCT03882437?cond=danon+disease&draw=2&rank=2>


# Natural History of Rapidly Progressing Heart Failure

## Cardiac Clinical Features

- Progressive hypertrophic cardiomyopathy/heart failure
- Key Clinical Biomarker Changes
  - Echo:
    - Worsening diastolic parameters
      - ↑ left ventricular end diastolic diameter (LVEDD)
      - ↓ left ventricular fractional shortening (LVFS)
      - ↑ ventricular wall thickness
      - ↓ left ventricular ejection fraction (LVEF) is late event
  - Hemodynamics: Decreasing cardiac output and/or stroke volume
  - Biomarkers: Elevated BNP, CK-MB, troponin

# Female Danon Cardiac Histology Suggests Broad LAMP2 Expression Important for Reversal of Phenotype

- Immunohistochemistry (IHC) from Danon female patients with severe disease display large patches negative for LAMP2 expression
- Broad expression of LAMP2 is likely the key to correcting phenotype rather than overall protein levels
- Based on this data, IHC demonstrating broad and *homogeneous cardiac expression may be the best predictor of long-term efficacy*



Cardiac IHC Staining in Female Danon Patient  
Requiring Transplant at 10 y<sup>1</sup>

# RP-A501: Subject Characteristics & AAV Vector Dose


| Patient ID | Age at Treatment | Dosing Weight | Cohort Dose                | Total Dose               |
|------------|------------------|---------------|----------------------------|--------------------------|
| 1001       | 17 y             | 52.2 kg       | $6.7 \times 10^{13}$ GC/kg | $3.25 \times 10^{15}$ GC |
| 1002       | 20 y             | 89.1 kg       | $6.7 \times 10^{13}$ GC/kg | $5.97 \times 10^{15}$ GC |
| 1005       | 18 y             | 97.8 kg       | $6.7 \times 10^{13}$ GC/kg | $6.08 \times 10^{15}$ GC |
| 1006       | 21 y             | 82.7 kg       | $1.1 \times 10^{14}$ GC/kg | $9.10 \times 10^{15}$ GC |
| 1007       | 20 y             | 96.7 kg       | $1.1 \times 10^{14}$ GC/kg | $1.06 \times 10^{16}$ GC |

# RP-A501 Demonstrated a Manageable Safety Profile

---

- In Low-dose cohort, RP-A501 was generally well tolerated with manageable safety profile
  - **Transient** and **reversible** decline in platelets
  - **Transient** and **reversible** transaminase elevation
- In Higher-dose cohort, a single patient experienced drug-related SAE related to complement activation
  - Patient with enhanced risk due to high weight & vector dose and pre-existing AAV immunity
  - Anticipated SAE of atypical hemolytic-uremic syndrome (aHUS) resulting in reversible thrombocytopenia and acute kidney injury (AKI)
  - AKI required supportive care including eculizumab and transient hemodialysis with full return to baseline kidney function within 2-3 weeks
- ***All patients have fully recovered from immune-related sequelae***

# RP-A501 Low Dose: DNA Vector Copy Number



\* Clinical course and VCN drop suggest apparent poor compliance with steroid regimen  
\*\* 1001, 1002 Month 12; 1005 Month 9




# RP-A501 Low Dose Cohort Demonstrates Robust Cardiac Expression as Measured by LAMP2 Immunohistochemistry (IHC)

| Patient | LAMP2B Relative Expression vs. Control* |        |                    |
|---------|-----------------------------------------|--------|--------------------|
|         | Regimen                                 | Week 8 | Long Term          |
| 1001    | Steroids only<br>(limited compliance)   | 7.8%   | <15%* <sup>1</sup> |
| 1002    | Steroids only<br>(local monitoring)     | 36.9%  | 67.8% <sup>1</sup> |
| 1005    | Steroids → Tacrolimus                   | 17.6%  | 92.4% <sup>2</sup> |


\* Endomyocardial biopsies were obtained and stained for LAMP2. Percent area of cell staining was quantitated using software in a blinded fashion and expression compared to normal heart tissue. Values represent average of 3-14 sections. Qualitative assessment reported for samples with high variance.

1. Sample obtained at Month 12
2. Sample obtained at Month 9


# RP-A501 Low Dose: Patients 1002 and 1005 Demonstrate Robust Cardiac Expression of LAMP2 by IHC Through Months 9 and 12, Respectively




**1002 Baseline**



**1002 Month 12**



**1005 Month 9**




**Normal Heart Control**

# RP-A501 Low Dose: Endocardial LAMP2B Protein Expression

| Patient | Relative LAMP2B Expression vs. Normal By Western Blot |                    |
|---------|-------------------------------------------------------|--------------------|
|         | Week 8                                                | Long Term          |
| 1001    | 20.7%                                                 | 17.9% <sup>1</sup> |
| 1002    | 27.3%                                                 | -                  |
| 1005    | 42.8%                                                 | 61.1% <sup>2</sup> |

1. Sample obtained at Month 6  
2. Sample obtained at Month 9


# RP-A501 Low Dose Improves or Stabilizes Key Cardiac Marker of Heart Failure: B-type Natriuretic Peptide (BNP)




# RP-A501 Electron Microscopy of Cardiac Myocytes Demonstrates Marked Decrease in Vacuolar Pathology: **Patient 1005**

---


Baseline



Week 8



Month 9



# RP-A501 Low Dose Confers Improvement in Cardiac Output Based on Invasive Hemodynamics in Patients 1002 and 1005

## Cardiac Output (L/min)

| Patient | Baseline | Long Term <sup>1</sup>             |
|---------|----------|------------------------------------|
| 1001    | 5.2      | 4.12 <sup>2</sup>                  |
| 1002    | 3.58     | 5.8 <sup>2</sup> (1.62x increase)  |
| 1005    | 4.5      | 6.08 <sup>3</sup> (1.35x increase) |

**1. Calculated Stroke Volume: 40% increase in Patient 1002 and 31% increase in Patient 1005**

2. Sample obtained at Month 12  
3. Sample obtained at Month 9

## RP-A501 Low Dose: Safety & Efficacy Findings (n=3)

---

- Generally, **well tolerated** with manageable safety profile in all low-dose patients
- **LAMP2B gene expression** demonstrated in cardiac biopsies from all patients
- **Enhanced cardiac expression** by IHC and Western blot in both patients whose compliance with transient immunosuppressive regimen was closely monitored
  - Consistent increases in percentage and level of IHC staining at later (9-12m) timepoints
- **Positive trends** in key biomarkers and efficacy endpoints
  - Qualitative improvement of vacuolar pathology
  - Clinical lab markers demonstrated improvement in patients 1002 and 1005
  - Trends towards stabilization and/or improvement in cardiac output
- **Benefit observed in all three patients** serves as clinical proof of concept as *Danon disease patients generally do not improve independently*

# Leukocyte Adhesion Deficiency-I (LAD-I)

## *Monogenic Immunodeficiency Disorder*

RP-L102  
Fanconi Anemia


RP-A501  
Danon Disease

RP-L201  
Leukocyte Adhesion Deficiency-I

RP-L301  
Pyruvate Kinase Deficiency

RP-L401  
Infantile Malignant Osteopetrosis

### OVERVIEW:



**Background:** Disorder characterized by recurring and ultimately fatal infections caused by *ITGB2* gene mutations

- >50% patients with severe variant: **60-75% mortality by age 2**



**Current Available Treatments:** Allogeneic hematopoietic stem cell transplant associated with significant graft failure and acute GVHD



**Addressable Market:** Estimated **25-50 pts** treatable per year for severe population; up to 100 for potential expansion into moderate population in the US + Europe with effective gene therapy



**RP-L201:** Preclinical studies show stable engraftment and phenotypic correction in murine models, with restored neutrophil migration capability



**Regulatory Designations:** Fast Track and Rare Pediatric Disease designations in the U.S.; Advanced Therapy Medicinal Product (ATMP) classification in EU; Orphan Drug designation in the U.S./EU

<sup>1</sup> Defective expression of  $\beta_2$  integrin on leukocytes limits their extravasation to inflamed sites.

# LAD-I Program Summary

## Ultra-rare Disease = Streamlined Regulatory Approach

### Potential design & clinical endpoints:

- **Target Patient Population:** Severe LAD-I patients (CD18<2%), ~2/3 mortality by 2y
- **Control:** Literature review of ~300 pts. (Rocket/academic collaborative publication<sup>1</sup>)
- **Potential Clinical Endpoints:** Modest correction of CD18 expression, survival

## Efficacy Trials & Registration Status – Ahead of Schedule

### Registration & study planning on-schedule:

- ✓ Orphan Drug (US/EU) and Pediatric Rare Disease (US) designations granted
- ✓ IND & Phase 1/2 cleared by FDA
- ✓ Spain IMPD cleared
- ✓ US PI (UCLA Dr. Don Kohn)
- ✓ Recruitment underway from around the globe
- 3 global sites planned in the US/EU

## Product/Manufacturing Optimization


### Process now optimized:

- ✓ VCN using GMP vector with transduction enhancers consistently ~3 (Target VCN >1)

<sup>1</sup>Almarza Novoa E, Kasbekar S, Thrasher AJ, Kohn DB, Sevilla J, Nguyen T, Schwartz JD, Bueren JA. Leukocyte adhesion deficiency-I: A comprehensive review of all published cases. J Allergy Clin Immunol Pract. 2018 Jan 20. pii: S2213-2198(17)31026-7. doi: 10.1016/j.jaip.2017.12.008.

# Rationale for Gene Therapy in LAD-I: CD18 Expression Correlates to Patient Survival

Kaplan-Meier Survival Estimates by Neutrophil CD18 Expression  
-Patients with moderate LAD-I not receiving allogeneic HSCT-




Natural history studies show the *correlation* between *higher CD18* expression and longer patient *survival*, supporting gene therapy’s potential in LAD-I patients

The grey diamond indicates the 39% survival to age 2 years for 66 evaluable patients with severe LAD-I not receiving HSCT

# LAD-I: Mouse Study Shows LAD-I Correction

- Primary and serially transplanted LAD mice underwent CD18 lenti GTx with different promoters
- Myeloablative conditioning was used
- Rocket chose the Chimeric cFES/CTSG (myeloid-specific) promoter (Post-transplant PB VCN 0.4-0.9)



# RP-L201 (LAD-I) Clinical Trial and Outcome Measures<sup>1</sup>

## Non-Randomized Phase 1/2 Study

### Design

- Enroll 9 pediatric patients globally
  - Phase 1: Enroll two patients to assess safety and tolerability
    - Enrollment Complete
  - Phase 2: Overall survival at multiple sites (US and Europe) n=7

### Primary Outcomes


- Phase 1:
  - Safety associated with treatment
- Phase 2:
  - Survival: proportion of patients alive at age 2 and at least 1-year post infusion (without HSCT)
  - Safety associated with treatment

### Secondary Outcomes

- Percentage of patients with at least 10% neutrophil CD18 expression
- Percentage of patients with at least 0.1 peripheral WBC gene marking (VCN) at 6 months post-infusion
- Incidence and severity of infections
- Improvement in neutrophils
- Resolution (partial or complete) of any underlying skin rash or periodontal abnormalities

<sup>1</sup>Source: <https://clinicaltrials.gov/ct2/show/NCT03812263?cond=Leukocyte+Adhesion+Deficiency&rank=5>

# Pre-Gene Therapy Medical History of Patient 1001




Historical patient records collected by UCLA Mattel Children's Hospital  
LAD has received CIRM Funding

# Patient 1001: 12-Month Follow-Up

## Key Drug Product Metrics


- CD34+ Cell Dose:  $4.2 \times 10^6$  cells/kg
- Drug Product VCN: 3.8

% CD18 Expression (PMN)



PMN: polymorphonuclear lymphocytes

VCN (PBMC)



PBMC: peripheral blood mononuclear cell

# Patient 1001: Visible Improvements Post-Treatment

**Pre GTx:** Severe infections  $\geq 1$  per year; numerous hospitalizations, severe skin lesions, continuous prophylactic antibiotics and required home schooling

**Post GTx:** No infections or hospitalizations, off antibiotics and able to attend school

## Spontaneous Abdominal Lesion



**Baseline  
(Pre-Treatment)**



**3-months  
(Post-Treatment)**




**6-months  
(Post-Treatment)**



**12-months  
(Post-Treatment)**

# Pre-Treatment Medical History of Patient 1004




# Patient 1004: 6-Month Follow-Up

## Key Drug Product Metrics

- CD34+ Cell Dose:  $2.8 \times 10^6$  cells/kg
- Drug Product VCN: 2.5


% CD18 Expression (PMN)



CD18 at baseline was reported as dim in approximately 63% PMNs, likely indicating an unstable protein, and in the context of additional clinical and laboratory evidence of severe LAD-I

PMN: polymorphonuclear lymphocytes

VCN (PBMC)




PBMC: peripheral blood mononuclear cell

# Medical History & Follow-up of Patient 2006

7-m.o. male

- Diagnosed at birth given family history of disease
- Delayed separation of umbilical cord (6 weeks)
- 2 older siblings diagnosed with severe LAD-I



## KEY DRUG PRODUCT METRICS

**CD34+ Cell Dose:**  
 $4.3 \times 10^6$  cells/kg

**Drug Product VCN:**  
2.87

Hematopoietic reconstitution observed Day 35 post-infusion

Clinically stable with no reported serious adverse events post-infusion

**76% CD18 expression** at 2-month timepoint

Historical patient records collected by UCLA Mattel Children's Hospital  
LAD has received CIRM Funding



# RP-L201 Study Summary

---

- Drug product produced in 7 out of 9 patients in Ph 1 & Ph 2
- Safety results of RP-L201 appear favorable
  - Infusion well tolerated; no drug product-related SAEs or severe AEs as of November 2020
- Preliminary Efficacy of CD18 PMN expression observed in both patients with  $\geq 6$ -months of follow-up
  - Patient 1001: durable CD18 PMN expression  $\sim 40\%$  and PB VCN of 1.2 at 12-months post-infusion and *resolution of skin lesions*
  - Patient 1004: CD18 PMN expression 23% 6-months post-treatment and PB VCN kinetics similar to those of first patient
  - Patient 2006: 2-months post-treatment had CD18 PMN expression of 76%
- Commercial-grade drug product and centralized testing for all patients treated
- Phase 2 study *enrollment and treatment expected to be completed 1H2021*

# Pyruvate Kinase Deficiency (PKD)

## Monogenic Red Blood Cell Hemolytic Disorder

RP-L102  
Fanconi Anemia


RP-A501  
Danon Disease

RP-L201  
Leukocyte Adhesion Deficiency-I

RP-L301  
Pyruvate Kinase Deficiency

RP-L401  
Infantile Malignant Osteopetrosis

### OVERVIEW:



**Current Available Treatments:** *Chronic* blood transfusions and splenectomy—side effects include iron overload and extensive *end-organ damage*



**Addressable Market<sup>2</sup>:** ~250-500 patients/year



- Conservative estimates conclude a number from 3,000 to 8,000 in the US + Europe combined



**RP-L301:** *Improvements in multiple disease components* in a PKD mouse model, including increased hemoglobin, reduced reticulocytosis, resolved splenomegaly and reduced hepatic erythroid clusters and iron deposits

**Regulatory Designations:** Fast Track in the US and Orphan Drug designation in the US/EU

<sup>1</sup>One glucose molecule is metabolized into two Phosphoenolpyruvate and ultimately two Pyruvate (pyruvic acid) molecules; this final enzymatic step yields two additional ATPs from each glucose molecule

<sup>2</sup>Market research indicates the application of gene therapy to broader populations could increase the annual market opportunity from approximately 250 to 500, based on an estimated prevalence in the US/EU of approximately 3,000 to 8,000.

# Preclinical Studies Demonstrated Safety and Efficacy of Lentiviral-mediated Gene Therapy

## PKD mice transplanted with gene-corrected cells demonstrated phenotypic correction:

- Significant increase in RBC count and half-life
- Decreased erythropoietin levels
- Normalized spleen and liver size & structure, with no evidence of erythroid clusters or iron deposits
- Improvement in red cell pyruvate kinase enzymatic pathway as assessed by metabolomic assays



## Favorable Safety Results:

- No physical, behavioral biochemical, hematologic or morphologic abnormalities observed in transplanted mice
- Limited evidence of PGK-coRPK-WPRE in nonhematopoietic organs, indicating very low risk of germline transmission
- No evidence of replication competent lentivirus (RCL)

# RP-L301: Global Phase 1 PKD Gene Therapy Study

## Primary Endpoint

Safety and toxicity of RP-L301

## Key Secondary Endpoints

- Clinically significant reduction of anemia
- **Transfusion independence** (when relevant) at 12-months
- Achievement of 50% reduction in transfusion requirements (when relevant) at 12-months
- **PB and BM** genetic correction as demonstrated by VCN
- Reduction of hemolysis

## Key Eligibility Criteria

### Inclusion:

- PKD diagnosis with a confirmed *PKLR* mutation
- Age:
  - 1<sup>st</sup> cohort (N=2): ≥18 to 50-years
  - 2<sup>nd</sup> cohort (N=2): ≥12 to 17-years
  - 3<sup>rd</sup> cohort (N=2): ≥ 8 to 11-years
- **Severe and/or transfusion-dependent anemia**
- Prior splenectomy
- Adequate cardiac, pulmonary, renal and hepatic function

### Clinical Sites:

- Hospital Universitario Fundación Jiménez Díaz, Madrid
- Stanford University, Palo Alto, California
- Hospital Infantil Universitario Niño Jesús, Madrid

# RP-L301: Patient Characteristics and Product Metrics

## Patient Characteristics


| Patient | Age (y) and Gender | Hemoglobin (g/dL) | Bilirubin (mg/dL) | Erythropoietin (mIU/mL) | Transfusion Requirement for 2 Years Prior to Enrollment |
|---------|--------------------|-------------------|-------------------|-------------------------|---------------------------------------------------------|
| 1001    | 31 F               | 7.4 <sup>†</sup>  | 13.4 mg/dL        | 35.6 mIU/mL             | ~14 transfusion episodes                                |
| 1002*   | 47 M               | 7.0 <sup>‡</sup>  | 7.4 mg/dL         | 57.2 mIU/mL             | ~5 transfusion episodes                                 |

## Product Metrics

| Patient | CD34+ Cells/kg        | Mean VCN: Liquid Culture |
|---------|-----------------------|--------------------------|
| 1001    | 3.9 x 10 <sup>6</sup> | 2.73                     |
| 1002*   | 2.4 x 10 <sup>6</sup> | 2.08                     |

\* Infused November 2020  
† Average hemoglobin calculated over 2-years prior to study enrollment  
‡ Average hemoglobin calculated over 2-years prior to study enrollment; patient has declined red blood cell transfusions

# RP-L301: Preliminary Efficacy Results—L301-006-1001




- Marked hemoglobin improvement ~7.4 g/dL to 14.3 g/dL
- No transfusion requirements following engraftment

\* Lab Values during mobilization/apheresis & post-conditioning period were not included

Data as of October 2020


# RP-L301: Preliminary Efficacy Results—Patient 1001

At 1–3 months post RP-L301 \*




Reticulocytes decreased

Erythropoietin normalized



Bilirubin decreased from 13.4 mg/dL to 1.8 mg/dL

Hepcidin increased from <4.0 ng/mL to 90.1 ng/mL



VCN in PBMCs 1.55 and VCN in BMMCs 2.72 at 3-months post RP-L301

\* Lab Values during mobilization/apheresis & post-conditioning period were not included  
Data as of October 2020

# RP-L301: Preliminary Safety Results

## Treatment-emergent Adverse Events (Grade 3 or higher) (N=1 patient)

| Event<br>System Organ Class (NCI CTCAE v. 5.0) | Adverse Events Grade |   |   |
|------------------------------------------------|----------------------|---|---|
|                                                | Any                  | 3 | 4 |
| <b>Blood and lymphatic system disorders</b>    |                      |   |   |
| Neutropenia                                    | 1                    | 1 | – |
| <b>Gastrointestinal disorders</b>              |                      |   |   |
| Stomatitis                                     | 1                    | 1 | – |
| <b>Investigations</b>                          |                      |   |   |
| AST increased                                  | 1                    | 1 | – |
| ALT increased                                  | 1                    | 1 | – |
| <b>Metabolism and nutrition</b>                |                      |   |   |
| Hypertriglyceridemia                           | 1                    | – | 1 |

- ***No RP-L301 related adverse events***
- ***Patient 1001 achieved neutrophil engraftment on day +13***

**Adverse events considered related to mobilization/apheresis (N=2 patients):**  
Grade 2 SAE (chest pain, dyspnea and nausea) during apheresis collection. These events were considered related to hyperleukocytosis and the mobilizing agents. They resolved with supportive care and without sequelae. Other events included Grade 2 bone pain and Grade 3 leukocytosis.

# RP-L301 Conclusion: Hemoglobin Normalized in First Patient

---

- Safety profile of RP-L301 *appears favorable*
  - Infusion well tolerated (n=1); no IP-related SAEs or AEs
  - Hematopoietic reconstitution in less than 2-weeks in initial patient
- Preliminary efficacy activity observed during initial 3-months after administration of RP-L301
  - Patient 1001 with peripheral blood VCN of 1.55 at 3-months, *hemoglobin nearly doubled* and normalized hemolysis markers (Hb from baseline *increased ~7g/dL* at 3-months post RP-L301)
- Second cohort will enroll older pediatric patients and is expected to be initiated in 1H2021

*Commercial-grade drug product and centralized testing for all treated patients*

# Infantile Malignant Osteopetrosis (IMO)

## *Monogenic bone resorption disorder*

RP-L102  
Fanconi Anemia

RP-A501  
Danon Disease

RP-L201  
Leukocyte Adhesion Deficiency-I

RP-L301  
Pyruvate Kinase Deficiency

RP-L401  
Infantile Malignant Osteopetrosis

### OVERVIEW:

-  **Background:** *Dysfunctional osteoclast* disease characterized by bone marrow failure, skeletal deformities, and neurologic abnormalities caused by *TCIRG1* mutations in >50% of cases<sup>1</sup>
  - *Frequent mortality in early years of life, severe marrow failure and visual impairment during 1<sup>st</sup> year*
-  **Current Available Treatments:** Hematopoietic stem cell transplants associated with GVHD and *limited efficacy*
-  **Addressable Market:** *>50* patients/year<sup>2</sup>
-  **RP-L401:** *In vitro* restoration of osteoclast resorptive function observed; *in vivo* correction in murine model
-  **Regulatory Designations:** Rare Pediatric Disease, Orphan Drug and Fast Track designations in the US


<sup>1</sup>Source: [https://www.orpha.net/consor/cgi-bin/OC\\_Exp.php?Ing=EN&Expert=667](https://www.orpha.net/consor/cgi-bin/OC_Exp.php?Ing=EN&Expert=667)

<sup>2</sup>Estimated incidence of one in 200,000 live births; Source: [http://www.orpha.net/consor/cgi-bin/OC\\_Exp.php?Ing=EN&Expert=667](http://www.orpha.net/consor/cgi-bin/OC_Exp.php?Ing=EN&Expert=667)


# Preclinical Mouse Data Supports Progression to Phase 1

Oc/oc mice receiving RP-L401 showed correction of the disease phenotype, with increased long-term survival, tooth eruption, weight gain, and normalized bone resorption

## Increased Long-term Survival



## Reversal of Osteopetrotic Bone Phenotype



# RP-L401 (IMO) Clinical Trial and Outcome Measures<sup>1</sup>

## Non- Randomized Phase 1 Study

### Design

- Enroll 2 patients, with a confirmed diagnosis of IMO with documented *TCIRG1* mutation
  - 1-month or older

### Primary Outcomes

- Safety associated with treatment

### Secondary Outcomes

- Normalization of serum calcium and blood counts
- Presence of gene-modified blood and bone marrow cells
- Normalization of bone abnormalities on X-ray and DEXA scans
- Prevention or stabilization of vision and hearing loss
- Reduction in hepatosplenomegaly

<sup>1</sup>Source: <https://www.clinicaltrials.gov/ct2/show/NCT04525352?term=NCT04525352&draw=2&rank=1>

# Growing IP Portfolio



## Multiple in-licensed patent families for GTx products and related technology platforms

### Supporting current pipeline efforts:

- Four In-licensed pending international patent applications filed under Patent Cooperation Treaty (PCT):
  - FA (2)
  - LAD-I
  - PKD
- Multiple patent applications pending:
  - Danon (exclusive world-wide license from UCSD)
- Multiple patent families licensed from REGENXBIO:
  - Danon – AAV9 (exclusive world-wide license)
  - Danon – 2 undisclosed capsid serotypes (exclusive world-wide option to license)
- Multiple cell and gene therapy platform technologies licensed for pipeline product improvements



## Rocket Proprietary Filed IP

### Extensive patent portfolio across multiple platforms:

- Multiple pending patent applications for ex-vivo LVV programs
- Multiple pending patent applications for in-vivo AAV

# World-Class Research and Development Partners



|                                               |                                        |                                     |                                       |
|-----------------------------------------------|----------------------------------------|-------------------------------------|---------------------------------------|
| CIBER                                         | IIS FJD                                | REGENXBIO                           | University of California, Los Angeles |
| CIEMAT                                        | Lund University                        | Stanford Medical School             | University of Minnesota               |
| Fred Hutchinson Cancer Research Center        | Memorial Sloan Kettering Cancer Center | UCL                                 | University of Pennsylvania            |
| Hospital Universitario Fundación Jiménez Díaz | Niño Jesús Hospital                    | University of California, San Diego |                                       |

# Expansion into Cranbury, NJ: R&D/CMC Efforts and Eventual cGMP Manufacturing

2021

- Continue R&D to *further support* CMC analytics and internal QC and release testing activities for RP-A501
- 50,000 sq. ft. from this facility will be *dedicated to AAV cGMP* manufacturing (FDA and EMA compliant)
- Initiate in-house GMP clinical manufacturing
- Enables *dual-sourcing* for Danon commercial capacity



RCKT Cranbury (NJ)  
103,720 sq. ft. production facility



# R&D & Manufacturing Facility



# R&D & Manufacturing Facility

---

# Near and Long-Term Value Drivers

*Potential for Five Gene Therapy Products to be Approved by 2025*

4Q2020

1H2021

2H2021

✓ LAD-I: Initiate Phase 2 Study

✓ LAD-I: Phase 1 Data Update

✓ IMO: Initiation of Clinical Study

✓ FA: Preliminary “Process B” Data

✓ PKD: Preliminary Phase 1 Data

✓ Danon: Preliminary Phase 1 Data

● FA: Updated “Process B” Data

● LAD-I: Initial Phase 2 Data

● PKD: Phase 1 Data Update

● IMO: Initial Phase 1 Data

● Danon: Updated Phase 1 Data