

Preliminary Conclusions Obtained in Fanconi Anemia Patients Treated by Lentiviral-mediated Gene Therapy after more than 2 years of Follow-up

Paula Río, PhD



paula.rio@ciemat.es

ASCGT 22nd Annual Meeting, Washington DC, April 29th 2019


DISCLOSURES:

- *The Innovative Therapies Div. at CIEMAT receives funding for FA gene therapy from Rocket Pharma*
 - *PGK-FANCA-Wpre* Lentiviral vector has been licensed to Rocket Pharma*
-

Hematopoietic Stem Cell Defect in Fanconi Anemia: Consequences for Gene Therapy




Ceccaldi et al., 2012



✓ Somatic Mosaicism: Natural Gene Therapy

(Waisfisz et al. Nat Genet 1999)


✓ Proliferative advantage of human FA-HSCs corrected by Gene Therapy (Rio*, Navarro* et al. Blood, 2017)



J.Surrallès

HSC collection and Gene Therapy Trials of FA-A patients

Fancostem



CD34⁺ selection



Fresh



Frozen

Inclusion criteria: One parameter higher than:

Hb: 8.0 g/dL; Neutrophils: 750/mm³; Platelets: 30,000/mm³.

CD34⁺



Pre-stimulation
(8 h)



PGKFANCA:LV (12h)



Final product
Transduced CD34⁺ cells

NO CONDITIONING



Fancolen

Inclusion criteria: One parameter lower than:

Hb: 8.0 g/dL; Neutrophils: 1,000/mm³; Platelets: 50,000/mm³.



FANCOSTEM: Purification of mPB CD34⁺ cells



	<i>Apheresis</i>			<i>CD34⁺ immunoselection</i>		
<i>Patient</i>	N ^{er} Aph.	<i>CD34⁺/kg</i>	<i>% CD34⁺</i>	<i>CD34⁺/kg</i>	<i>% CD34⁺</i>	<i>Recovery</i>
01003	3	1.6x10 ⁶	0.05	8.6x10 ⁵	17.4	53.2
02002	2	7.2x10 ⁶	0.16	5.1x10 ⁶	80.0	70.9
02003	3	7.6x10 ⁶	0.17	2.9x10 ⁶	29.3	47.8
02004	3	6.6x10 ⁶	0.15	2.0x10 ⁶	25.6	26.6
02005	2	4.8x10 ⁶	0.11	1.1x10 ⁶	48.7	22.7
01005	2	2.6x10 ⁶	0.18	1.0x10 ⁶	51.8	48.7
02006 *	2	3.1x10 ⁶	0.12	1.3x10 ⁶	8.4*	41.0
02007 *	3	9.9x10 ⁵	0.03	6.0x10 ⁵	1.8*	60.6
02008 \$	2	2.8x10 ⁶	0.08	3.4x10 ⁵	1.5\$	12.0
MEDIAN (Range)	2 (2-3)	3.1x10⁶ (0.9-7.6x10 ⁶)	0.12 (0.03-0.18)	1.1x10⁶ (0.6-5.1x10 ⁶)	25.6 (1.8-80.0)	48.0 (22.7-70.9)

* CD34⁺ cell immunoselection using depletion tubing set



\$ CD34⁺ cell immunoselection with gentle washing

Strategy to mobilize and transduce HSCs from FA-A patients included in FANCOLEN



FA-02005 (4 years old): Hb 12.5 g/dL; NEUTROPHILS 1,680/mm³; **PLATELETS 38,000/mm³**

FA-02006 (6.6 years old): Hb 11.3 g/dL; **NEUTROPHILS 760/mm³**; PLATELETS 79,000/mm³



FA-02002 (5 years old): Hb 10.5 g/dL; NEUTROPHILS 1,600/mm³; **PLATELETS 29,000/mm³**




FA-02004 (7.6 years old): Hb 10.8 g/dL; NEUTROPHILS 900/mm³; **PLATELETS 46,000/mm³**

FANCOLEN: Infusion products



<i>PATIENT</i>	<i>N^{er} of infused cells/kg</i>	<i>CD34⁺ cells/kg</i>	<i>% CD34⁺</i>	<i>Mean VCN/cell (CFCs)</i>	<i>Transduced CD34⁺cells/kg</i>	<i>Transduced CFCs/kg</i>	<i>MMC survival</i>
<i>02005 (Fresh)</i>	1.7x10 ⁶	1.40x10 ⁶	80.5%	0.17	2.3x10 ⁵	2.8x10 ³	12.1%
<i>02002 (Cryopr)</i>	6.0x10 ⁵	0.56x10 ⁶	90.6%	0.45	2.5x10 ⁵	1.7x10 ⁴	30.3%
<i>02004 (Cryopr)</i>	1.2x10 ⁶	0.72x10 ⁶	57.0%	0.23	1.7x10 ⁵	6.9x10 ³	27.0%
<i>02006* (Fresh)</i>	2.1x10 ⁷	0.77x10 ⁶	3.5%	0.53	4.1x10 ⁵	1.6 x10 ⁵	49.2%





Kinetics of Gene-marked Cells in PB of Non-conditioned FA Patients Infused with Transduced CD34⁺ cells



Additional preliminary data from 24-36 month follow-up indicate progressive increases in PB VCNs





Not analyzed

Progressive increase of gene-marked subpopulations in PB



Not analyzed



Progressive increase of gene-marked subpopulations in BM including CD34⁺ cells



Not analyzed

Integration Site Analysis in Patient 02002

% corr PBCs: 0.03 0.27 0.14 N.D 2.1 6.5 9.2 12.1 16.5 22.1 34.4 42.5 54.9





Wang, Wei (GeneWerk GmbH)





Session: Clinical Gene Therapies for Blood Diseases

April 29th, 11:15-11:30 (Location: Jefferson)

Progressive Phenotypic correction of BM CFCs (MMC-Resistance)




Progressive Phenotypic correction of PB T Cells (DEB-Induced Chromosomal Instability)




Kinetics of corrected and uncorrected PB leukocytes prior to and after Gene Therapy


FA-02002








FA-02006



FA-02005



FA-02004






---●--- Uncorrected leukocytes/ μ L




---●--- Corrected leukocytes/ μ L

PB Cell Counts in FA Patients prior to and after Gene Therapy




02002 (Cryo)
 2.5×10^5 cCD34⁺/Kg
 1.7×10^4 cCFU/Kg






02006 (Fresh)
 4.0×10^5 cCD34⁺/Kg
 1.6×10^5 cCFU/Kg



02005 (Fresh)
 2.3×10^5 cCD34⁺/Kg
 2.8×10^3 cCFU/Kg



02004 (Cryo)
 1.7×10^5 cCD34⁺/Kg
 6.9×10^3 cCFU/Kg



Months after gene therapy

Months after gene therapy

Months after gene therapy

Conclusions

- ❑ No Severe Adverse Events
- ❑ Collection of clinically relevant numbers of mPB CD34⁺ cells
- ❑ Infusion of corrected CD34⁺ cells in non-conditioned FA patients:
 - ✓ Engraftment and repopulation advantage of gene-corrected HSPCs
 - ✓ Oligoclonal reconstitution and gene correction of multipotent HSCs
 - ✓ Phenotypic correction of BM progenitors and PB T cells
 - ✓ Stabilization of the BMF
 - ✓ Additional preliminary data from 24-36 month follow-up indicate progressive genetic and phenotypic reversal

Next Steps



- *FANCOLEN I: Long term follow up protocol (15 years)*
- *FANCOLEN II: Hospital Niño Jesús (Madrid)*
 - Infusion of higher numbers of corrected HSCs:
 - *Collection / infusion of HSCs prior to BMF (Fresh/Fresh)*
 - *Improved transduction of HSCs (transduction enhancers)*
 - Commercial vector
- *Stanford University: Phase 1 Clinical Trial (n=2)*